

## [<sup>35</sup>S]-amino acids

|                 |                                   |   |
|-----------------|-----------------------------------|---|
| <sup>35</sup> S | L-Methionine, [ <sup>35</sup> S]- | CH <sub>3</sub> S(CH <sub>2</sub> ) <sub>2</sub> CH(NH <sub>2</sub> )COOH;<br>In 50 mM Tris-HCl, pH 7.4 containing<br>10 mM 2-mercaptoethanol;<br>> 1000 Ci/mmol; 10-20 mCi/ml  |
|                 | Met-[ <sup>35</sup> S]-label      | L-Methionine, [ <sup>35</sup> S], L-Cysteine, [ <sup>35</sup> S];<br>E. coli hydrolysate containing 70% L-<br>Methionine, [ <sup>35</sup> S]-, ~25% L-Cysteine, [ <sup>35</sup> S]-<br>and ~5% miscellaneous amino acids;<br>> 1000 Ci/mmol |
|                 | L-Cystein, [ <sup>35</sup> S]-    | HSCH <sub>2</sub> CH(NH <sub>2</sub> )COOH;<br>In stabilized aqueous solution containing<br>10mM DTT;<br>>1000 Ci/mmol; 10 mCi/ml   |

## [<sup>35</sup>S]-nucleotide analogue

|                 |  |   |
|-----------------|--|---|
| <sup>35</sup> S | <b>3'-Phosphoadenosine-5'-phosphosulfate (PAPS), [<sup>35</sup>S]-</b> | In aqueous solution containing 50%<br>ethanol;<br>>1000 Ci/mmol; 1 mCi/ml |
|-----------------|--|---|

You need different concentrations or specific activities. Please send us your request.